



Regulatory T cells in draining lymph nodes of *Lawsonia intracellularis* infection in pigs

Riber, Ulla; Andreassen, Elisa W.; Jungersen, Gregers

Publication date:
2013

[Link back to DTU Orbit](#)

Citation (APA):

Riber, U., Andreassen, E. W., & Jungersen, G. (2013). *Regulatory T cells in draining lymph nodes of Lawsonia intracellularis infection in pigs*. Abstract from 10th International Veterinary Immunology Symposium, Milano, Italy.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

10th International Veterinary Immunology Symposium – IVIS 2013, Milan Italy

Abstract (page 104), Poster

Ulla Riber, Elisa W. Andreassen, Gregers Jungersen

National Veterinary Institute, Technical University of Denmark

Regulatory T cells in draining lymph nodes of *Lawsonia intracellularis* infection in pigs

Lawsonia intracellularis infection in pigs cause diarrhoea and poor performance in growing pigs and is an important contributor to the high antibiotic usage in pig production. Experimentally, a primary subclinical *L. intracellularis* infection can induce protection against a secondary challenge infection. Although, immune responses to *L. intracellularis* infection have been investigated to a certain level, with IFN- γ being a key factor for development of protection, the role of T_{regs} is unknown. Activation of suppressive T_{regs} may play a role in the ability of *L. intracellularis* to survive in the infected host.

Four pigs were challenged twice with *L. intracellularis* infectious material, with four weeks interval. Lack of faecal shedding after the second challenge indicated the pigs were protected. The pigs developed *L. intracellularis* specific IgG responses and CMI responses in PBMCs confirmed T_C cells (CD3⁺CD4⁻CD8 β ⁺) and memory T_H cells (CD3⁺CD4⁺CD8 α ⁺) being main producers of IFN- γ . Pigs were slaughtered 8 week after the second challenge and ileocacal lymph node cells (iLNC) and PBMCs were prepared and frozen.

With focus on identification and characterisation of T_{regs}, iLNC were co-cultured with porcine IL-2 and *L. intracellularis* antigen (Ag), Con A, or IL-2 alone. Before culture iLNC showed 1.4-4.0% Tregs (CD3⁺FoxP3⁺), which were mainly CD25^h. ILNCs were around 20% CD4⁺CD8 α ⁺ T cells of which 6.3-10.7% were T_{regs}, whereas within CD4⁺CD8 α ⁻ T cells (37%) and CD4⁻CD8 α ⁺ T cells (35%) the levels of T_{regs} were 1.7-3.4% and 0.9-1.6%, respectively. The phenotype CD4⁺CD8 α ⁺ of T_{regs} may indicate these cells being induced (iT_{regs}) compared to naturally occurring (nT_{regs}) mainly CD4⁺CD8 α ⁻.

Co-culture for 6 days (CFSE proliferation assay) with IL-2 and Con A identified FoxP3⁺ cells among proliferating cells, however proliferation in Ag-cultures was at same level as without antigen.

Further characterisation of T_{regs} after *L. intracellularis* antigen culture of iLNC and PBMC will be performed.